

Remarks concerning an entropy-like quantity S_q especially for quantum systems with a particle picture; model with pure point spectra^(*)

S. SCHLIEDER

*Max-Planck-Institut für Physik, Werner-Heisenberg Institut
Föhringer Ring 6, D-80805 Munich, Germany*

(ricevuto il 27 Gennaio 1998; approvato il 15 Ottobre 1999)

Summary. — For quantum-mechanical systems an entropy-like quantity S_q is defined. S_q can differ from the usually defined entropy S and S_q may increase with time for an isolated system. The essential condition for the difference between S and S_q is the assumption that the set \mathbf{A} of observables which can be represented by a measurement is a proper subset of the set of self-adjoint operators. The underlying idea is made visible in the case of particle systems with non-trivial scattering. The model character of the reasoning comes from the fact that continuous spectra are replaced by point spectra. So it seems evident that no direct connection exists between S_q and the Sinai-Kolmogorov entropy at least in this model with pure point spectra.

PACS 03.65.Bz – Foundations, theory of measurement, miscellaneous theories (including Aharonov-Bohm effect, Bell inequalities, Berry's phase).

1. – Introduction

Thermodynamics and the physical quantities appearing in it got a deeper foundation by that branch of theoretical physics, which one calls “Statistical Mechanics” and which is connected with the work of Boltzmann and Gibbs. Using statistical methods the concepts of atomism of matter were introduced into the framework of the older thermodynamics. This way also the thermodynamic quantity “Entropy” got a new interpretation. Entropy from that time on expressed the deficit of knowledge about the exact details of the atomistic picture of a physical system—for instance for a gas the lack of knowledge concerning the position or the momentum of each particle as an individuum in the ensemble representing this gas.

(*) The author of this paper has agreed to not receive the proofs for correction.

This statistical approach is then continued if one proceeds from classical systems to quantum systems. Since one cannot characterize the state of the system by using position and momentum of every particle at the same time, one has to go over to the quantum state of the entire system into which for instance—in the case of a sufficient dynamical separation of each particle from the other ones—the quantum state of each particle is an ingredient. The lack of knowledge will be described by a set of probabilities $\{w_i\}$ for the possible quantum states $\{\varphi_i\}$ respectively and one defines the entropy S by

$$(1.1) \quad S = - \sum w_i \ln w_i, \quad \sum w_i = 1.$$

If one regards entropy in this way, defined as a function of time, then an irritating feature may arise in the cases where the $\{\varphi_i\}$ do not describe stationary states. Let $\{\varphi_i\}$ be an orthonormal basis in the Hilbert space \mathbf{H} with the interpretation $\{\varphi_i = \varphi_i(t_0)\}$ and $\{\varphi_i(t) = U(t, t_0)\varphi_i(t_0)\}$ describing the unitary time development of the Schrödinger states $\{\varphi_i(t)\}$. The entropy itself however remains constant, since the orthonormal basis $\{\varphi_i(0)\}$ transforms under the time development into the one parameter set $\{\varphi_i(t)\}$ of orthonormal bases; this means that the mixture $\{w_i\varphi_i(0)\}$ transforms into the mixtures $\{w_i\varphi_i(t)\}$; so $S(t) = S(0)$ and the entropy remains constant.

In one respect this feature is not a new experience: In the classical cases, one defines entropy at a time t for instance at $t = 0$ usually by the support $\Pi(0)$ of the probability measure at $t = 0$ in the phase space. Generically the support $\Pi(0)$ changes with time to supports $\Pi(t)$; however as a consequence of one of the famous Liouville theorems, the volume of Π stays constant and thereby also the entropy connected with this volume remains also constant.

On the other side, entropy—as a quantitative expression for the lack of knowledge about the system—should generically increase with time. Also one can learn in thermodynamics that the entropy of an isolated system should increase with time except in the special cases, where it has reached its internal equilibrium.

These controversial features have been very well known for a long time and as a consequence new aspects were brought into the game to overcome these difficulties. So one introduced besides the fine grain picture in phase space a coarse grain picture by performing local smoothing of the probability measure. Another approach to these problems is possible since Kolmogorov's fundamental work. One uses an adequate decomposition of the phase space to define a certain kind of entropy—conditioned by the knowledge of the past (the so-called “Kolmogorov-Sinai Entropy”). This may increase in time with a specific rate for certain dynamical systems.

The following considerations are restricted to quantum-mechanical systems and should lead to the definition of an entropy-like quantity S_q . Usually the entropy concept for quantum-mechanical systems is based on the uncertainty in the knowledge about the quantum states actually involved. Typically one expresses this fact with the use of mixtures of pure states.

In contrast to the procedure described defining the entropy for a quantum-mechanical system we intend to take into account also the question whether all virtual information contained in a quantum state can be made into real information by experiments. The generalized answer to this question leads to the definition of an entropy-like quantity S_q in sect. 3. A general feature of S_q is that it can be $\neq 0$ also for pure states and that S_q can also grow with time for one and the same quantum state. To restrict the discussion to this main point, we specialize the discussion to states which are pure before

measurements are performed. The increase of S_q with time is a common feature with the increase of the Kolmogorov-Sinai entropy. However the reasons for the increase are quite different; one sees this immediately regarding the fact that here a model with pure point spectra is discussed.

In short, the definition of S_q is based on the following consideration: If one tries to describe reality in a physical theory, it is not completely clear with which objects of the theory reality is coupled [1]. The structure of quantum mechanics remains unchanged in the following. However the concept of a quantum state is looked upon as not immediately related to physical reality. The relation to it is more intimately given by the results of measurements on the physical systems being in a certain quantum state. The author has heard of this approach the first time from W. Thirring [2]. On the other hand, the preference for the concept of a quantum state to represent the results of measurements (as special types of events)—instead of describing an object—is also in a certain concordance with Haag’s opinion [3] about the meaning of the concept of a quantum state.

In the spirit of the remarks made before one is led to Shannon’s concept of entropy which was carried further by Kolmogorov and Khinchin [4]. To demonstrate this, one can start with a “scheme” (the word used by Khinchin), containing n events. In a trial exactly one of the events a_1, a_2, \dots, a_n can take place with the probabilities w_1, w_2, \dots , respectively w_n , and with $\sum w_i = 1$: The “schema” is written as

$$(1.2) \quad \begin{pmatrix} a_1 a_2 \cdots a_n \\ w_1 w_2 \cdots w_n \end{pmatrix} \quad \text{with} \quad \sum w_i = 1.$$

One defines the entropy of such a scheme—one can also call this an n -alternative with a probability measure—by the well-known expression

$$(1.3) \quad S = - \sum w_i \ln w_i.$$

The maximum of S is assumed with $w_i = 1/n$ and has the value $S = \ln n$. The minimum $S = 0$ is reached, if one of the events, say a_k , takes place with certainty $w_k = 1$; the scheme then becomes trivial.

The concept of entropy of that kind can be used in two ways:

– Given a schema the above-defined entropy is an uncertainty in the answer for the question, which event actually will happen, when the scheme will be realized in an experiment.

– After the outcome of the experiment is known and in this way the uncertainty is replaced by certainty the amount of information gained can also be expressed by the defined entropy.

In the quantum-mechanical case alternatives with a probability measure originate for instance in connection with the measurements of observables. The connection appears most clearly in cases where the observables can be represented by self-adjoint operators with pure point spectra. Without special assumptions one cannot expect that such interesting quantities for instance as the momentum or the energy of a quantum system have as their representatives self-adjoint operators with pure point spectra. On the other hand, there are known ways—partly of mathematical nature, partly by somewhat changing the physical situation—to replace the continua in the spectra by point spectra. In this case—assuming in addition that the point spectrum is simple, *i.e.* to each eigenvalue

corresponds a one-dimensional subspace, the corresponding scheme for the quantum state φ and the observable A describes the situation

$$(1.4) \quad \begin{pmatrix} a_1, a_2, \dots, a_k, \dots \\ w_1, w_2, \dots, w_k, \dots \end{pmatrix}.$$

The interpretation for this scheme is

$$(1.5) \quad \varphi = \sum c_i \varphi_i, \{A\varphi_i = a_i \varphi_i\}, \{w_i = |c_i|^2\}.$$

The corresponding entropy $\tilde{S}(\varphi, A)$ is then

$$(1.6) \quad \tilde{S}(\varphi, A) = - \sum w_i \ln w_i,$$

\tilde{S} is the expression for the uncertainty in the result measuring the observable A on the system in the state φ before the measurement is made.

Section 2 contains some simple remarks concerning some relations between different $\tilde{S}(\varphi, A_j)$ by fixing φ and varying A_j .

The concept of alternatives with probabilities seems to be an adequate tool for the definition of an entropy-like quantity, if one regards events as the primary concepts linking the theory to the physical reality. On the other hand, one should clarify the relationship between quantities like $\tilde{S}(\varphi, A_j)$ and the entropy S (defined in the usual way) of a system which is based only on the uncertainty concerning the quantum states. $\tilde{S}(\varphi, A_j)$ is quantitatively equal to the entropy of the mixture which originates from the measurement of A_j . But what does this have in common with the usual $S(\varphi)$, which in our case, treated first for simplicity for $\varphi \in H$, vanishes? In sect. 3 we intend by using expressions like $\tilde{S}(\varphi, A_j)$ to come back to an entropy-like quantity $S(\varphi)$. However, there remains a conceptional difference between $S(\varphi)$ and $S_q(\varphi)$. $S(\varphi)$ is based on the more ontological concept of a state, while $S_q(\varphi)$ expresses the knowledge concerning the probabilities of possible events concentrated in the concept of a state. Therefore $S(\varphi)$ and $S_q(\varphi)$ may differ from each other.

In sect. 4 one finds some remarks concerning S_q for particle systems and in sect. 5 especially for 2-particle systems. In sect. 6 a critical discussion and some concluding remarks are added.

2. – Alternatives weighted with probability measures (schemes) and their entropies

Two schemes M_1 and M_2 may be given. Certain conditions may exist between the probabilities of the events of M_1 and those of M_2 . We are interested in the simplest situations, where one scheme is finer (or coarser) than the other. Let us assume M_1 is finer than M_2 . For the members of M_1 we use double indices:

$$(2.1) \quad M_1 : \begin{pmatrix} a_{11}, a_{12}, \dots, a_{1r}, a_{21}, a_{22}, \dots, a_{2s}, \dots, a_{m1}, a_{m2}, \dots, a_{mz} \\ w_{11}, w_{12}, \dots, w_{1r}, w_{21}, w_{22}, \dots, w_{2s}, \dots, w_{m1}, w_{m2}, \dots, w_{mz} \end{pmatrix},$$

$$(2.2) \quad M_2 : \begin{pmatrix} b_1, b_2, \dots, b_m \\ w_1, w_2, \dots, w_m \end{pmatrix}.$$

If one of the events in scheme M_1 , namely a_{11} or a_{12} or a_{1r} happens, then in M_2 event b_1 takes place, if a_{21} or a_{22} or a_{2s} in scheme M_1 happens, then b_2 in M_2, \dots and so on. For the probabilities one assumes

$$(2.3) \quad \sum_k w_{rk} = w_r, \quad \sum_r w_r = 1, \quad r = 1, 2, \dots, m.$$

In this case, where M_1 is finer than M_2 , one can write

$$(2.4) \quad M_1 \stackrel{f}{\supset} M_2.$$

It is clear, how one can use the property of being finer or coarser for observables especially here for those with pure point spectra. Besides the common domain of definition the corresponding self-adjoint operators must commute and the property of being finer and coarser should be independent of the state to which they are applied. If these conditions are fulfilled, one can introduce sequences of self-adjoint operators

$$(2.5) \quad A_1 \stackrel{f}{\subset} A_2 \stackrel{f}{\subset} \dots \stackrel{f}{\subset} A_n,$$

where the operators become finer going from left to right. In principle they can end to the right with a self-adjoint operator with a simple point spectrum. As one can see easily, the corresponding entropies increase also in the direction to the right

$$(2.6) \quad \tilde{S}(\varphi, A_1) \leq \tilde{S}(\varphi, A_2) \leq \dots \leq \tilde{S}(\varphi, A_n).$$

If as assumed, φ is a pure state, then $\tilde{S}(\varphi, 1) = 0$, where 1 is the unit-operator.

That such a sequence becomes arbitrarily fine and ends with a self-adjoint operator with a simple point spectrum can be easily fulfilled in a separable Hilbert space. In the following sections it is however intended to restrict the observables to a set which is represented by a proper subset of the set of self-adjoint operators. Therefore the following assumption is not trivial and generically an idealisation.

Assumption (F): Each observable represented by a self-adjoint operator with a pure point spectrum is an element of a sequence of observables which ends—in the direction to become finer—with an observable having a simple point spectrum.

To sum up we arrive at the following situation:

1) The interesting observables from a physical standpoint are approximated by observables, whose representing self-adjoint operators have pure point spectra.

2) Further one assumes that in this approximating set one can find self-adjoint operators to fulfill (F).

Two remarks should be helpful:

1) An important motivation of this paper is the fact that not all self-adjoint operators (respectively, those with pure point spectra) are actual observables.

2) Generically one can derive other observables from the spectrum of some observable or from the spectra of different observables—if the corresponding self-adjoint operators commute—by using real functions of the values contained in their spectra. For instance, for a system with several particles the total momentum or the total energy is calculated

from values of the momenta and energies measured for each particle separately. This procedure must be justified by the assumption that the S -operator exists and therefore the dynamical interactions between the particles can be neglected. So without the intention to discriminate such observables as the total momentum, total energy in the case of the particle systems we have in mind the special observables belonging to the measurements made at each particle. When we refer in the following to the set of observables in the particle picture we restrict this set in such a way that the measurements of the physical quantities of each particle are represented separately. To give a name to this set one can call it: "Set of actual observables in the particle picture". To give a general description of the set of actual observables, one can remark that they are those which actually transform by their measurement generically a pure state into a mixture or a mixture into a more refined mixture⁽¹⁾.

3. – Definition of the entropy-like quantity S_q for quantum-mechanical systems

Although we will use the concept of the entropy-like quantity S_q only for quantum systems consisting of several particles, we will describe the general situation, in which S_q may differ from S for quantum states.

As mentioned before, the concept of a quantum state will here not be looked upon as describing an ontological object but as a tool to describe all possible events in itself or originating from interaction with other systems and specially to make propositions concerning the probabilities of the results of measurements in the future.

For the definition of $S_q(\varphi)$ we intend therefore to make use of the quantities $\tilde{S}(\varphi, A_j)$ with variable A_j . One can differentiate between two properties of the quantities $\tilde{S}(\varphi, A_j)$.

1) If one performs a preparation of a state φ as an individual of an ensemble, then this procedure is intended to create a uniform ensemble with all individuals in the same state, or in the language of statistics, to create an ensemble with the highest possible order. In a certain idealisation the outcome of the procedure will then be a pure state $\varphi \in H$. Generically the measurement of an observable is connected with a disturbance of this order (expressed by the transformation of the pure state φ to a mixture). In order to define $S_q(\varphi)$ one is inclined to use those A_j in $\tilde{S}(\varphi, A_j)$ which lead to the smallest disturbance of this order.

2) If, on the other hand, one regards a sequence of observables as in (2.5) in the direction to the left, where the observables become coarser, one could for instance end the procedure with the unit-operator. The effect is then: No disturbance, however also no information. In the whole, one has the feature that coarser operators lead to smaller entropies $\tilde{S}(\varphi, A_j)$. However, one is interested in gaining as much information about the state as possible at all.

Combining 1) and 2) in pursuing the intention to define $S_q(\varphi)$ by quantities $\tilde{S}(\varphi, A_j)$ one has to look for those A_j leading to the highest possible information, together with the least disturbance of the quantum state in question.

From remark 2), it follows that only observables A_j with the finest properties are used for the definition of $S_q(\varphi)$ by the expressions $\tilde{S}(\varphi, A_j)$. In this context we use

⁽¹⁾ The remark 2) has its origin in a discussion with H. Roos who criticized the too narrow concept of observables in an earlier version of this paper.

the assumption (F) made in sect. 2. Let us use the notation \mathbf{A}_e for the set of actual observables with simple point spectra.

Definition: $S_q(\varphi) = \inf_{A_k \in \mathbf{A}_e} \tilde{S}(\varphi, A_k)$.

Remark 1: Let us denote the set of self-adjoint operators with simple point spectrum by $\hat{\mathbf{A}}_e$. In the case that $\mathbf{A}_e = \hat{\mathbf{A}}_e$ for the quantum-mechanical system in question, one gets $S_q(\varphi) = 0$. We assumed that φ is a pure state and we can find self-adjoint operators in $\hat{\mathbf{A}}_e$ with simple point spectra for which φ is an eigenstate. In these cases—even if one includes mixtures besides the pure states—the S_q defined above is equal to entropy S as usually defined.

Remark 2: The sets \mathbf{A}_e , respectively $\hat{\mathbf{A}}_e$, can be replaced by the sets \mathbf{P}_e , respectively $\hat{\mathbf{P}}_e$, of the minimal projection operators belonging to the spectral decompositions of the self-adjoint operators in \mathbf{A}_e , respectively $\hat{\mathbf{A}}_e$.

The main point of this section is clear from the foregoing: To define S_q one has to define the set \mathbf{A}_e of all simple actual observables as a subset of all simple self-adjoint operators $\hat{\mathbf{A}}_e$ (with pure point spectra). If $\mathbf{A}_e = \hat{\mathbf{A}}_e$ is a good idealisation for the states of a Hilbert space \mathbf{H} (with a certain physical interpretation), then $S_q(\phi)$ is the same quantity as $S(\phi)$, where ϕ is a pure state or a mixture. In the next section we consider a different physical situation.

Remark 3: The definition of $S_q(\varphi)$ is based on $\tilde{S}(\varphi, A_k)$ with $A_k \in \mathbf{A}_e$ and also on condition (F) which clearly has its origin in the model used. Physical considerations make it desirable to have self-adjoint observables with the highest distinguishing power in $\tilde{S}(\varphi, A_n)$. Therefore one can think for instance of introducing maximal Abelian subalgebras of self-adjoint operators instead of \mathbf{A}_e . This idea was also discussed with other physicists. However, the degree of fineness of the operators involved in the definition of S_q —on which for instance the amount of entropy creation in a scattering process depends—has its limitations in physical circumstances and is not so much dependent on the mathematical tools one uses.

4. – Actual observables for a physical system of n -particles

The essential concept expressed by the definition of the entropy-like quantity S_q shall be illustrated with the example of a system consisting of n particles. The Hilbert space \mathbf{H} of the total system is constructed from the 1-particle Hilbert spaces $H^{(k)}$ with $k = 1, 2, \dots, n$ by taking the n -fold tensor product and performing after that the completion. If the particles are of the same kind one has to restrict \mathbf{H} by symmetry (or antisymmetry) conditions applied to the combinations of the products of the 1-particle states. For the physical conditions we have in mind that

- 1) The energy of the particles is low enough to exclude particle creation.
- 2) The particles behave like free particles apart from the short time when they are scattering off each other.

So one can use the mathematical construction of the tensor product of 1-particle states as a description of the physical situation of the n -particle state in a good approximation.

It is clear what physical conditions are necessary for that: The n -particle state has to be dilute enough that the time of scattering should be short compared with the time

when the particles move approximately as free particles. This is a stronger assumption than that of the existence of the S -matrix.

Under these conditions one is able to define the set of observables in a plausible way in view of the physics involved.

The set of the actual observables is a subset of the set

$$(4.1) \quad \mathbf{A} = \{A_j^{(1)} \otimes A_k^{(2)} \otimes \cdots \otimes A_r^{(n)}\}.$$

Thereby the observables with different upper indices operate in different 1-particle spaces. To avoid any confusion in the registration of the measurement results (which result belongs to which particle?) or to be free from correlations arising from the interactions of measuring devices among themselves, it might sometimes be necessary to restrict the regions in which the observables in (4.1) actually operate. However, this shall not be discussed here in detail; only the meaning of the word subset of the set (4.1) should be made plausible.

The set \mathbf{A} of (4.1) representing the actual observables belongs only to a small subset of the self-adjoint operators operating in \mathbf{H} . The observables of (4.1) have a characteristic property: They transform superpositions of factorizing states, which usually are the outcome of scattering processes into the corresponding mixtures, if they have been measured.

As follows from the content of sect. 3, one has to use proper subsets of the 1-particle operators denoted by $\{A_j^{(l)}\}$ $l = 1, 2, \dots, n$ in (4.1) for the definition of $S_q(\Phi)$, $\Phi \in \mathbf{H}$. The subsets are collections of those observables which are as fine as possible—in the model used here they are represented by self-adjoint operators with a simple (or non-degenerate) point spectrum. If one uses the symbolic notation \mathbf{A}_e for the set $\mathbf{A}_e^{(1)} \otimes \mathbf{A}_e^{(2)} \otimes \cdots \otimes \mathbf{A}_e^{(n)}$, the definition for S_q is then

$$(4.2) \quad S_q(\Phi) = \inf_{\Phi \in \mathbf{H}, C_j \in \mathbf{A}_e} \tilde{S}(\Phi, C_j).$$

The measurements represented by an element of \mathbf{A}_e (this means by the tensor product of n self adjoint operators operating in the n 1-particle spaces) transforms $\Phi \in \mathbf{H}$ into the following mixture:

$$(4.3) \quad \Phi = \sum_{j_1, j_2, \dots, j_n} \varphi_{j_1} \otimes \cdots \otimes \varphi_{j_n} \longrightarrow \{|c_{j_1, j_2, \dots, j_n}|^2 \varphi_{j_1} \otimes \varphi_{j_2} \otimes \cdots \otimes \varphi_{j_n}\}$$

with

$$(4.4) \quad \sum_{j_1, j_2, \dots, j_n} |c_{j_1, j_2, \dots, j_n}|^2 = 1$$

if $\|\Phi\| = 1$. Thereby the orthonormal basis $\{\varphi_{j_1}\}, \{\varphi_{j_2}\}, \dots, \{\varphi_{j_n}\}$ chosen in each 1-particle space is the set of eigenstates of the corresponding self-adjoint operators:

$$(4.5) \quad A \in \mathbf{A}_e, A = A^{(1)} \otimes A^{(2)} \otimes \cdots \otimes A^{(n)}, \{A^{(l)} \varphi_{l_k} = a_{l_k} \varphi_{l_k}\}$$

with $l = 1, 2, \dots, n$.

Obviously one gets

$$(4.6) \quad \tilde{S}(\Phi, A) = - \sum_{j_1, j_2, \dots, j_n} |c_{j_1, j_2, \dots, j_n}|^2 \ln |c_{j_1, j_2, \dots, j_n}|^2$$

and one should vary $A \in \mathbf{A}_e$, defined above to obtain

$$(4.7) \quad S_q(\Phi) = \inf_{A_j \in \mathbf{A}_e} \tilde{S}(\Phi, A_j).$$

It is a trivial remark that $S(\Phi) = 0$ is valid independently of the kind of scattering or whether there is any scattering at all.

However, by restricting the actual observables by (4.1) and basing the entropy-like quantity S_q on the expressions $\tilde{S}(\Phi, A_j)$, one gets another picture for S_q . Considering a single scattering process described by $\Phi(t)$ (the mathematical details of the asymptotic conditions should not matter in the moment), and looking to $S_q(\Phi(t))$ at a time t_{in} before and a time t_{out} after the scattering took place one has the following generic picture.

Before the scattering of 2 particles one prepares each particle if possible as a pure 1-particle state (or at least as a well-defined mixture of 1-particle states). So one has before scattering a state $\varphi^{(1)} \otimes \chi^{(2)}$ as a tensor product (or, taking statistics into account, a symmetrized or antisymmetrized tensor product if one has particles of the same kind).

Denoting $\Phi_{\text{in}} = \varphi_{\text{in}}^{(1)} \otimes \chi_{\text{in}}^{(2)}$ the state before the scattering one can find $A^{(1)}$, respectively $B_{\text{in}}^{(2)}$, for which $\varphi_{\text{in}}^{(1)}$, respectively $\chi_{\text{in}}^{(2)}$, are eigenstates of $A^{(1)}$, respectively $B^{(2)}$.

Therefore $S_q(\Phi_{\text{in}}) = \inf \tilde{S}(\Phi_{\text{in}}, A_j) = 0$ with $A_j \in \mathbf{A}_e$. However, after the scattering process the state Φ_{out} does not factorize. Therefore it is not possible to find an operator $B \in \mathbf{A}_e$, for which Φ_{out} is an eigenstate; this has the consequence $S_q(\Phi_{\text{out}}) > 0$. The change of a factorizing state into a non-factorizing state by a scattering process is clearly a fundamental feature for quantum-mechanical scattering processes. If one would try to describe a non-factorizing state in the particle picture, which means by 1-particle properties, this would not be successful; this can only be done after transforming the quantum state—here the state Φ_{out} —into a mixture by performing a measurement corresponding to an operator of the kind defined in (4.1). In this sense the quantities $\tilde{S}(\Phi, A_j)$ are used as quantized expressions for the uncertainty in the results of measurements before the measurements are performed—the uncertainty concerns here the question, which factorizing state would come out in a measuring process corresponding to (4.3). One can find some discussion in the concluding remarks of sect. 6 concerning the mathematical frame using in principle n measurements performed for some n -particle state—leading to a description which does not correspond to the physical situation, if the number n of particles is large compared to 2.

5. – Some remarks concerning S_q for 2-particle states

The content of this section is the discussion of the quantities S_q in the case of 2-particle quantum states. We use here the Schrödinger picture. It is a trivial consequence of the discussion in sect. 3 and sect. 4 that one is not able to define S_q as a characteristic quantity for a Heisenberg state.

The state $\Phi(t_{\text{in}}) = \Phi_{\text{in}}$ before the scattering is transformed by the scattering process into $\Phi(t_{\text{out}}) = \Phi_{\text{out}}$. For the considerations here it is not important whether one is able to describe the scattering by a unitary time development $\Phi(t) = U(t, t_{\text{in}})\Phi_{\text{in}}$ coming to

$\Phi_{\text{out}} = U(t_{\text{out}}, t_{\text{in}})\Phi_{\text{in}}$ or to come from Φ_{in} immediately to Φ_{out} by applying to Φ_{in} a scattering operator (S -matrix). In a certain idealization the preparation of Φ_{in} might lead to

$$(5.1) \quad \Phi_{\text{in}} = \psi_{\text{in}}^{(1)} \otimes \chi_{\text{in}}^{(2)},$$

with

$$(5.2) \quad \psi_{\text{in}}^{(1)} \in \mathbf{H}^{(1)}, \chi_{\text{in}}^{(2)} \in \mathbf{H}^{(2)}, \|\psi_{\text{in}}^{(1)}\| = 1, \|\chi_{\text{in}}^{(2)}\| = 1.$$

(For reasons of simplicity it is assumed here that the particles are of different type, so it is not necessary to symmetrize or antisymmetrize.) One sees that for

$$(5.3) \quad S_q(\Phi_{\text{in}}) = \inf_{A_j \in \mathbf{A}_e^{(1)}, B_k \in \mathbf{A}_e^{(2)}} \tilde{S}(\Phi_{\text{in}}, A_j \otimes B_k),$$

$$(5.4) \quad S_q(\Phi_{\text{in}}) = 0.$$

The property that $S_q(\Phi_{\text{in}})$ vanishes is based on the fact that one can find in $\mathbf{A}_e^{(1)}$ as well as in $\mathbf{A}_e^{(2)}$ observables, which are represented by self-adjoint operators with simple point spectra and for which $\psi_{\text{in}}^{(1)}$, respectively $\psi_{\text{in}}^{(2)}$, are eigenstates. Φ_{out} can be described by

$$(5.5) \quad \Phi_{\text{out}} = \sum_{i,k} c_{ik} \hat{\varphi}_i \otimes \hat{\eta}_k, \quad \sum_{i,k} |c_{ik}|^2 = 1$$

with $\{\hat{\varphi}_i\}$, respectively $\{\hat{\eta}_k\}$, as orthonormal systems in $\mathbf{H}^{(1)}$, respectively $\mathbf{H}^{(2)}$.

J. v. Neumann has shown that one can find orthonormal systems $\{\varphi_l\}$, respectively $\{\eta_k\}$ (dependent on Φ), to bring Φ , here in particular Φ_{out} , into the normal form

$$(5.6) \quad \Phi_{\text{out}} = \sum_e \sqrt{w_l} \varphi_l \otimes \eta_l$$

with $\sum w_l = 1$.

If in the sum of (5.6) some w_i coincide, one gets a variety of normal forms for one and the same state. If for instance for a certain state Φ one has a total degeneration of the w_i , one gets

$$(5.7) \quad \Phi = \frac{1}{\sqrt{n}} \sum_{l=1}^n \varphi_l \otimes \eta_l = \frac{1}{\sqrt{n}} \sum_{l=1}^n \varphi'_l \otimes \eta'_l$$

with

$$(5.8) \quad \varphi_l = \sum_{s=1}^n u_{sl} \varphi'_s, \quad \eta_l = \sum_{s=1}^n \bar{u}_{sl} \eta'_s.$$

Thereby $\{u_{sl}\}$ is an arbitrary unitary $n \times n$ -matrix and $\{\bar{u}_{sl}\}$ is its complex conjugate. The normal forms of the state Φ_{out} show in their structure the correlations originating from the conservation of physical quantities like total momentum or total energy. The

point spectrum would physically correspond to the enclosure of the 2-particle system in a box. (Such an idealization comes, however, somewhat in conflict with the S -matrix picture.)

On the other hand, the normal forms give a hint which pairs of observables one should use as factors in the tensor product $A_f \otimes B_g$ to obtain the infimum, which here becomes the minimum. Before demonstrating this, it is useful to give a

Definition:

A state $\Phi \in \mathbf{H}$, for instance Φ_{out} in (5.1), may be represented in a normal form

$$(5.9) \quad \Phi = \sum \sqrt{w_l} \varphi_l \otimes \eta_l.$$

As a second mathematical object there should be given a pair of self-adjoint operators with pure simple point spectra and their tensor product $A \otimes B$. We say $A \otimes B$ is adapted to the given normal form of Φ if $\{\varphi_l\}$, respectively $\{\eta_l\}$, is in the set of eigenvectors of A , respectively B .

With this definition one is able to formulate the

Proposition: The quantity

$$(5.10) \quad \inf_{C_j \in \mathbf{A}_e^{(1)}, D_k \in \mathbf{A}_e^{(2)}} \tilde{S}(\Phi, C_j \otimes D_k)$$

is reached by a pair $A \otimes B$, which is adapted to one and the same normal form of Φ .

Remark 1: The infimum is actually a minimum and one gets

$$(5.11) \quad S_q(\Phi) = \min \tilde{S}(\Phi, C_j \otimes D_k) = \tilde{S}(\Phi, A \otimes B) = - \sum_l w_l \ln w_l.$$

Remark 2: If there exist several normal forms for Φ and the tensor product $A' \otimes B'$, with $A' \in \mathbf{A}_e^{(1)}$, $B' \in \mathbf{A}_e^{(2)}$ is adapted to another normal form of Φ , then again one gets

$$(5.12) \quad S_q(\Phi) = \tilde{S}(\Phi, A' \otimes B') = - \sum_l w_l \ln w_l.$$

Proof: We assume that both factors in $A \otimes B$ are adapted to the same normal form of Φ . $C \otimes D$ should be constructed with arbitrary operators C and D , fulfilling the conditions $A, C \in \mathbf{A}_e^{(1)}$ and $B, D \in \mathbf{A}_e^{(2)}$. The proposition has been proven if one can show the validity of the following inequalities, respectively equalities,

$$(5.13) \quad \tilde{S}(\Phi, C \otimes D) \geq \tilde{S}(\Phi, C \otimes 1) \geq \tilde{S}(\Phi, A \otimes 1) = \tilde{S}(\Phi, A \otimes B).$$

We prove the different parts of (5.13):

$\tilde{S}(\Phi, C \otimes D) \geq \tilde{S}(\Phi, C \otimes 1)$ holds, because $C \otimes D$ is finer than $C \otimes 1$. In addition $\tilde{S}(\Phi, A \otimes 1) = \tilde{S}(\Phi, A \otimes B)$, since $A \otimes B$ is adapted to one and the same normal form of Φ . With $\Phi = \sum \sqrt{w_l} \varphi_l \otimes \eta_l$ this is true, since

$$(5.14) \quad (P_k \otimes 1)\Phi = (P_k \otimes Q_k)\Phi = (1 \otimes Q_k)\Phi = \sqrt{w_k} \varphi_k \otimes \eta_k$$

can be derived from the properties $P_k \varphi_l = \delta_{kl} \varphi_l$, $Q_k \eta_l = \delta_{kl} \eta_l$.
The remaining task is to prove

$$(5.15) \quad \tilde{S}(\Phi, C \otimes 1) \geq \tilde{S}(\Phi, A \otimes 1) = - \sum w_l \ln w_l.$$

Let now $\{\hat{\varphi}_s\}$ be an orthonormal basis of eigenvectors of C . We use the substitution

$$(5.16) \quad \left\{ \varphi_l = \sum_s u_{sl} \hat{\varphi}_s \right\}$$

with $\{u_{sl}\}$ as unitary matrix. Introducing this substitution into (5.2) one gets

$$(5.17) \quad \Phi = \sum_l \sqrt{w_l} \varphi_l \otimes \eta_l = \sum_l \sqrt{w_l} \sum_s u_{sl} \hat{\varphi}_s \otimes \eta_l$$

and with $\hat{P}_s \hat{\varphi}_t = \delta_{st} \hat{\varphi}_t$

$$(5.18) \quad \begin{aligned} \|\hat{P}_s \otimes 1 \Phi\|^2 &= (\Phi, \hat{P}_s \otimes 1 \Phi) \\ &= \left(\sum_l \sqrt{w_l} \sum_t u_{tl} \hat{\varphi}_t \otimes \eta_l, \sum_m \sqrt{w_m} u_{sm} \hat{\varphi}_s \otimes \eta_m \right) \\ &= \sum_l w_l |u_{sl}|^2. \end{aligned}$$

So it follows

$$(5.19) \quad \tilde{S}(\Phi, C \otimes 1) = - \sum_s \sum_l w_l |u_{sl}|^2 \ln \sum_m w_m |u_{sm}|^2.$$

Since $f(x) = x \ln x$ is a convex function

$$(5.20) \quad f\left(\sum_l p_l x_l\right) \leq \sum_l p_l f(x_l)$$

provided

$$(5.21) \quad 0 \leq p_l \leq 1, l = 1, 2, \dots, \sum p_l = 1.$$

Introducing $|u_{sl}|^2 = p_l$, $w_l = x_l$ the inequality

$$(5.22) \quad \sum_l w_l |u_{sl}|^2 \ln \sum_m w_m |u_{sm}|^2 \leq \sum_l |u_{sl}|^2 w_l \ln w_l$$

holds for each s . Therefore one gets

$$(5.23) \quad \sum_s \sum_l w_l |u_{sl}|^2 \ln \sum_m w_m |u_{sm}|^2 \leq \sum_l w_l \ln w_l$$

and from (5.7)

$$(5.24) \quad \tilde{S}(\Phi, C \otimes 1) \geq - \sum_l w_l \ln w_l = \tilde{S}(\Phi, A \otimes B).$$

This completes the proof.

One may ask the question in which situation the equation

$$(5.25) \quad \tilde{S}(\Phi, A \otimes 1) = \tilde{S}(\Phi, C \otimes 1)$$

holds.

We discuss at first the case when all w_k are different from each other. There is a possibility that in the substitution for fixed s a certain k exists, that $|u_{sk}|^2 = 1$, while $|u_{sl}|^2 = 0$ for $l \neq k$. If this same property is true for every s , then the equation $\tilde{S}(\Phi, A \otimes 1) = \tilde{S}(\Phi, C \otimes 1)$ is valid. On the other hand, the unitary matrix (u_{sk}) in the substitution has only the effect to permute the eigenstates.

A bit more complicated is the case that some (or all) of the w_k are equal. For the corresponding step in the proof, where the convex function $f(x)$ is involved, one has then to take this function several times for the same values of the arguments by introducing the different values $x = w_l$ into the convexity condition. Clearly the convexity condition is also valid for this case. On the other hand, one is able to arrive at other normal forms for Φ by using unitary substitutions in the subspaces with equal values w_l . Then clearly one has a greater variety, besides A also other $C_j \in \mathbf{A}_e^{(1)}$, if they leave these subspaces invariant. Then $\tilde{S}(\Phi, A \otimes 1) = \tilde{S}(\Phi, C_j \otimes 1)$ follows. If one takes into account that one wants also $\tilde{S}(\Phi, C_j \otimes 1) = \tilde{S}(\Phi, C_j \otimes D_k)$, it is clear that again $C_j \otimes D_k$ must be adapted to a new normal form, which one obtains from unitary substitutions characterized above in (5.3). This means—when C_j is constructed by a unitary substitution (u_{sl}) from A as characterized above— D_k must be obtained by the unitary substitution (\bar{u}_{ls}) from B .

6. – Conclusions and some hints for a further approach

If the actual observables are only a proper subset of the self-adjoint operators, then the above-defined entropy-like quantity S_q differs from the usual entropy S . $S_q(\varphi)$ can be different from zero also for pure states φ and $S_q(\varphi)$ can grow with time; this is a special feature of scattering processes. We showed the mechanism, which leads to an increase of S_q for the 2-particle system without using the details of the scattering. This was demonstrated by using von Neumann's standard forms for the state after scattering. To make the physical picture consistent one should show by further considerations that the different components of the normal form belonging to different factorizing states can really be observed independently. That is a tedious discussion—as is often the case when physical consideration must be introduced into the mathematical framework—which nearly always contains elements of idealisation.

On the other hand, the idealisation introduced above for the characterization of the set of actual observables $\mathbf{A}_l = A_e^{(1)} \otimes A_e^{(2)} \cdots \otimes A_e^{(n)}$ becomes very idealistic, if n is large and if it is taken seriously as an expression for a measuring operation. In that case it is natural to use another picture—2-particle scattering states on a background consisting of the $(n-2)$ particle-system—, leading to a steady production of the entropy-like quantity S_q . This is a feature which S_q has in common with the Kolmogorov-Sinai

entropy, although the entropy production mechanism seems to be different. I think it is not worthless to study the entropy-production of S_q for particle systems with large n , using the considerations referring to the 2-particle scattering states.

* * *

I want to thank W. THIRRING for a discussion and his hint concerning the concept of Kolmogorov-Sinai entropy. Further it is a pleasure and a duty to mention colleagues at the Max Planck Institute, P. BREITENLOHNER, W. ZIMMERMANN, D. MAISON and E. SEILER.

REFERENCES

- [1] LUDWIG G., *Deutung des Begriffes physikalische Theorie und axiomatische Grundlegung der Hilbertraumstruktur der Quantenmechanik durch Hauptsätze des Messens* (Springer-Verlag) 1970.
- [2] THIRRING W., in numerous talks and papers.
- [3] HAAG R., in numerous talks and papers.
- [4] KHINCHIN A. I., *Mathematical Foundations of Information Theory* (Dover Publications, New York) 1957.